



***Agropyron pinifolium* Nevski (Poaceae): a new species record for the flora of Turkey**

Evren CABİ^{*1}, Ersin KARABACAK², Burçin ÇİNGAY³

¹ Department of Biology, Namık Kemal University, 59030, Tekirdağ, Turkey

² Department of Biology, 18 Mart University Çanakkale Botanic Garden, 17020, Çanakkale, Turkey

³ Nezahat Gökyiğit Botanic Garden, İstanbul, Turkey

Abstract

An addition to the known Turkish grasses, *Agropyron pinifolium* Nevski (Poaceae) is recorded for the first time from Turkey. This new record is confined to Vize province in Kırklareli. An updated description and notes on ecology and phenology of the new record are also presented. In addition, distribution map and illustration of this new record from Turkey is given.

Key words: Poaceae, *Agropyron pinifolium* Nevski, new record, Turkey

----- * -----

***Agropyron pinifolium* Nevski (Poaceae): Türkiye florası için yeni bir tür kaydı**

Özet

Bilinen Türkiye Buğdaygillerine ek olarak, *Agropyron pinifolium* Nevski (Poaceae) türü Türkiye'den ilk defa kaydedildi. Bu yeni kayıt Kırklareli ilinin Vize ilçesinde sınırlanmaktadır. Güncel bir betim ve bunun yanında türün ekolojisi ve fenolojisi üzerine notlar sunuldu. Ek olarak bu türe ait bir çizim ve dağılım haritası verildi.

Anahtar kelimeler: Buğdaygiller, *Agropyron pinifolium* Nevski,, yeni kayıt, Türkiye

1. Introduction

The taxonomy of *Agropyron* Gaertn. has been studied in different ways by various researchers. In a broad sense it was once thought to be one of the largest genera in the Triticeae, encompassing more than 100 species (Dewey, 1983). Nevski (1934) restricted *Agropyron* to perennial taxa with keeled glumes, a group of species referred to in English as the Crested Wheat grasses. He placed the other taxa in *Elytrigia* Desv, *Roegneria* C.Koch and *Elymus* L.. Subsequent works showed that members of *Agropyron* sensu Nevski are diploids, tetraploids, or hexaploids in which only the **P** genome is present) (Dewey and Asay, 1975; Dewey, 1983; Melderis, 1978; Assadi, 1995; Jensen et al., 2006; Genome designations as recommended by the International Triticeae Consortium; <http://herbarium.usu.edu/Triticeae/genmsymb.htm>). This narrow concept of *Agropyron* is now followed in most taxonomic works (e.g., Tzvelev, 1976; Melderis et al., 1980; Clayton and Renvoize, 1986; Wu et al., 2006; Barkworth et al., 2007; Cabi and Doğan, 2012). It is also supported by intergeneric crossing experiments (Assadi and Runemark, 1995). Much confusion prevails in regard to the number of species in *Agropyron*.

In the Flora of Turkey, Melderis (1985) recognized only one species, *A. cristatum* s.l. in *Agropyron* and subdivided it into two subspecies (subsp. *incanum* and subsp. *pectinatum*). He stated that subsp. *incanum* was confined to the high mountain steppes of East Anatolia whereas subsp. *pectinatum* grew throughout Turkey. *Agropyron cristatum* subsp. *pectinatum* was further divided into two varieties, one of which was var. *imbricatum* having pilose spikelets and the other one var. *pectinatum* having glabrous spikelets (Melderis, 1985).

Löve (1984) recognized another species from Turkey that he named *A. deweyi*. The seeds of this species were collected by J.R. Harlan in 1948 and cultivated in Evans Farm, Utah, U.S.A. He noted that this species might be a

* Corresponding author / Haberleşmeden sorumlu yazar: Tel.: +902822502670; Fax.: +902822502670; E-mail: ecabi2004@yahoo.com

variant of *A. cristatum* that arose under cultivated conditions far from its native habitat. Baum et al. neotypified the species in 2008.

The authors carried out intensive field studies between 2010 and 2014 and collected huge amount of herbarium specimens of the tribe Poeae R.Br. Population size, phenological traits and ecological preferences were observed in the field (Davis and Heywood, 1973). During these field studies particular attention was paid to a *Agropyron* population collected from edge of Vize to Kıyıköy (A1 Kırklareli, A1 sensu Davis, 1965) in 2013. Upon closer examination and going through Flora of Turkey (Davis, 1985) and other relevant floras, such as Flora Orientalis (Boissier, 1884), Flora of Syria, Palestine and Sinai (Post, 1933), Flora of Iraq (Bor, 1968), Flora Iranica (Bor, 1970), and Grasses of the Soviet Union (Tzvelev, 1983), it was identified as *A. pinifolium*. This species was only known from Crimea and Caucasus (From Novorossiisk to Tuapse) so far.

2. Materials and methods

The specimens were collected from Kırklareli: Vize to Kıyıköy, about 5 km from Vize, calcareous stony places, 41°59'25.7" N 27°8'22.73" E, 400 m, 16 June 2013, E.Cabi & E.Karabacak (Figure 3). Related herbarium specimens were studied in ANK, GAZI, ISTE, ISTF, VANF, E, K, and BM herbarium.

3. Results

Taxonomy

AGROPYRON Gaertn., *Novi Comment. Acad. Sci. Imp. Petrop.* 14(1): 539 (1770). / **OTLAK AYRIĞI**

Syn. Agropyrum Roem. & Schult., *Syst. Veg.* 2: 750 (1817). *Australopyrum* (Tzvelev) Á.Löve, *Feddes Repert.* 95(7-8): 442 (1984).

Lectotype: *A. cristatum* (L.) Gaertn., To Nash: "N.L. Britton & A. Brown, *Ill. Fl. N.U.S.* ed. 2. 1: 283 (1913)".

Agropyron pinifolium Nevski, *Trudy Sredne-Aziatsk. Gosud. Univ.*, Ser. 8b, Bot. 17: 57 1934. / **İbrelî ayrık** (Figure 1).

Syn.: *Agropyron sclerophyllum* (Novopokr.) Novopokr. *Sci. J. Rostov-on-Don State Univ.* 6: 39 1935. *Agropyron cristatum* subsp. *sclerophyllum* Novopokr. *Uchen. Zap. Rostovsk. Univ. V. M. Molotova, Trudy Biol. Fak.* 6: 39 1935.

A. karadaghense Kotov *Journ. Bot. Acad. Sci. Ukraine*, 5(1): 32. 1948. *Agropyron cristatum* subsp. *pinifolium* (Nevski) Bondar ex Korovina *Byull Vses Ord. Lenina Inst. Rast. N.I. Vavilov*, 81: 35. 1978.

Description

Caespitose perennial, Flowering shoots 15-24 cm. Sterile shoots forms dense mats with stems more or less thickened at base, and with numerous shortened sterile shoots up to 5 cm with distichously arranged, short, convolute and bent leaves. Cauline leaves linear, convolute. Spike up to 2.5 cm long and 1.4 cm wide, dense, rachis segments are visible. Spikelets 3-5 flowered. Glumes subequal, ovate-lanceolate with prominent keel and awn up to 4 mm long, tangled ciliate at keel. Lemma 5.5-6.5 cm long with an awn up to 4 mm long, glabrescent or scabrous. Palea nearly as long as the lemma, bidentate at the summit, sparsely ciliate on the keels (Figure 2).

Described from the vicinity of Gelendzhik. Type in Leningrad.

Flowers in May - July. Anemophilous. 2n=?

Specimens examined. Kırklareli: Vize to Kıyıköy, about 5 km from Vize, calcareous stony places, 41°59'25.7" N 27°8'22.73" E, 400 m, 16 June 2013, E.Cabi & E.Karabacak (Figure 3).

Habitat and elevation in Turkey - Calcareous stony places, woodlands' openings, 400 m. Flowering. May to July. It grows together with the following taxa *Stipa* sp., *Galium* sp., *Salvia* sp., *Poa* sp., *Catapodium* sp., *Bromus* sp. and *Aegilops* sp..

Distribution outside Turkey -West Transcaucasia, Crimea.



Figure 1. *Agropyron pinifolium* (=ibreli ayrık) in its habitat




Figure 2. *Agropyron pinifolium* (=ibreli ayrık). **A**, a general view of the species; **B**, spike; **C**, leaf detail; **D**, glume; **E**, lemma; **F**, palea. Illustrated by E.Karabacak from materials at NAKU and CBB.



Figure 3. The distribution map of *Agropyron pinifolium* (●) in Turkey

Use and economic value. - Forage grass.

Intensive studies in literature, herbaria and databases for identifying the specimens have been revealed that the identification is correct and it is a new record for the Flora of Turkey.

The herbarium specimens of *A. pinifolium* from Turkey are deposited at the herbaria of NAKU, CBB, and NGBB.

Acknowledgements

Thanks are due to the curators of herbaria ANK, GAZI, ISTE, ISTF, VANF, E, K, and BM, who allowed us to study their *Agropyron* specimens, and to the Namık Kemal University (NKUBAP.00.10.AR.12.10) for their financial assistance for the study.

References

- Assadi, M., Runemark H. 1995. Hybridization, genomic constitution and generic delimitation in *Elymus* s.l. (Poaceae, Triticeae). *Plant Systematics and Evolution* 194:189–205.
- Assadi, M. 1995. Meiotic configuration and chromosome number in some Iranian species of *Elymus*. *Botanical Journal of Linnean Society* 117:159–168.
- Barkworth M.E., Campbell J.J.N., Salomon, B. 2007. *Agropyron*. In: Barkworth, M.E., Capels, K.M., Long, S., Anderton, L.K. & Piep, M.B. (eds.) *Flora of North America*, Vol. 24, Poaceae, part 1, pp. 277–279. New York: Oxford University Press.
- Baum, B.R., Yen, C., & Yang, J.L. 2008. Neotypification of *Agropyron deweyi* (Poaceae, Triticeae). *Taxon* 18: 415–417.
- Boissier, P.E., 1884. *Flora Orientalis* 5, 749, Basileae, Geneva & Lugduni.
- Bor, N.L., 1968. Gramineae, in Townsend, C. C., Guest E., Al-Rawi (ed), *Flora of Iraq*, vol. 9. Ministry of Agriculture, Baghdad.
- Bor, N.L., 1970. Gramineae, in Rechinger, K. H. (ed), *Flora Iranica*, Vol. 70/30. Graz, Austria: Akademische Druk-Und Verlagsanstalt, Wien.
- Cabi, E., Doğan, M. 2012. Poaceae.: Güner, A., Aslan, S., Ekim, T., Vural, M. & Babaç, M.T. (eds.). *Türkiye Bitkileri Listesi (Damarlı Bitkiler)*. Ali Nihat Gökyiğit (ANG) Vakfı Yayın No 1, Flora Araştırmaları Derneği Yayın No: 1, İstanbul.
- Clayton, S.D., Renvoize, S.A. 1986. *Genera Graminum: Grasses of the world*. *Kew Bulletin. Additional Series* 13: 1–389.
- Davis, P.H., Heywood, V.H. 1973. *Principles of Angiosperm Taxonomy*. Huntington, New York: Robert E. Kieger Publishing Co.
- Davis, P.H. (ed.) 1965. *Flora of Turkey and East Aegean Islands*, Vol. 1, Edinburgh Univ. Press., Edinburgh.
- Davis, P.H. (ed.) 1985. *Flora of Turkey and East Aegean Islands*, Vol. 9, Edinburgh Univ. Press., Edinburgh.
- Dewey, D.R., Asay, K.H. 1975. The crested wheatgrasses of Iran. *Crop Science* 15:844–849.
- Dewey, D.R. 1983. Historical and current taxonomic perspectives of *Agropyron*, *Elymus*, and related genera. *Crop Sci.* 23:637–642.
- Jensen, K.B., Larson, S.R., Waldron, B.L., Asay, K.H. 2006. Cytogenetic and molecular characterization of hybrids between 6x, 4x, and 2x ploidy levels in crested wheatgrass. *Crop Science* 46:105–112.
- Knowles, R.P. 1955. A study of variability in crested wheatgrass. *Canadian Journal of Botany* 33: 534–546.
- Löve, A. 1984. *Conspectus of the Triticeae*. *Feddes Repertorium*. 95: 425–521.
- Melderis, A. 1978. Taxonomic notes on the tribe Triticeae (Gramineae) with special reference to the genera *Elymus* L. s.l. and *Agropyron* Gaertner s.l. *Botanical Journal of Linnean Society* 76: 369–384.
- Melderis, A. 1985 *Agropyron* (Gaertner). In: Davis, P.H. (ed.) *Flora of Turkey and the East Aegean Islands*, Vol. 9, pp. 204–206. Edinburgh Univ. Press., Edinburgh.
- Melderis, A, Humpries, C.J., Tutin, T.G., Heathcote, S.A. (1980). Tribe Triticeae Dumort. In: Tutin, T.G., Heywood V.H., Burges N.A., Moore, D.M., Valentine, D.H., Walters, S.M., Webb, D.A. (eds). *Flora Europaea*, Vol. 5. pp. 190–200. Cambridge University Press, Cambridge, England.
- Nevski, S.A. 1934. Tribe XIV. Hordeae Benth. In: Komorov V.L. *Flora of the U.S.S.R.* pp. 590–722. The Botanical Institute of the Academy of Sciences of the USSR. Leningrad, USSR.
- Post, G.E., 1933. *Flora of Syria, Palestine and Sinai*. (ed. 2, revised and enlarged by J.E. Dinsmore) Beirut: American Press.
- Tzvelev, N.N. 1983. *Grasses of the Soviet Union*, vol. 1-2. New Delhi: Amerind Publishing Company].
- Tzvelev, N.N. 1976. Poaceae URSS. Tribe 3. Triticeae Dumort. Genus 17. *Agropyron*. pp. 143–150. U.S.S.R Academy of Science Press, Leningrad.
- Wu, Z.Y., Raven, P.H., Hong, D.Y. 2006. (Eds.). *Flora of China*. Vol 22. Poaceae. Science press, Beijing and Missouri Botanical Garden Press, St. Louis.

(Received for publication 20 October 2014; The date of publication 15 April 2015)